

## Author index

- Abe, T. *see* Fukushima, S.  
 Agrén, M. *see* Brismar, H.  
 Agrén, M. *see* Ibarra, F.R.  
 Åhrén, B. *see* Persson, K.  
 Aizman, O. *see* Ibarra, F.R.  
 Aleksandrova, N.P., Goloubeva, E.V. & Isaev, G.G.  
 Mechanisms of genioglossus responses to inspiratory resistive load in rabbits, 253  
 Allescher, H.D. *see* Storr, M.  
 Amici, D. *see* Bramucci, M.  
 Aoki, K. *see* Kondo, N.  
 Aperia, A. *see* Ibarra, F.R.  
 Arvola, L., Hassaf, D., Melnikov, A.L., Helgesen, K.G. & Ytrehus, K. Adenosine induces prolonged anti- $\beta$ -adrenergic effects in guinea-pig papillary muscle, 11  
 Bentzen, J.G. *see* Sjaastad, I.  
 Bergström, G. *see* Nyström, H.C.  
 Berthon, P.M. *see* Lemoine, S.  
 Bigard, X. *see* Joumaa, W.H.  
 Björnström, K., Sjölander, A., Schipert, Å. & Eintrei, C. A tyrosine kinase regulates propofol-induced modulation of the  $\beta$ -subunit of the GABA<sub>A</sub> receptor and release of intracellular calcium in cortical rat neurones, 227  
 Bolton, T.B. *see* Kadinov, B.  
 Bramucci, M., Miano, A., Gobetti, A., Zerani, M., Quasinti, L., Maccari, E., Murri, O. & Amici, D. Bradykinin is not involved in angiotensin converting enzyme modulation of ovarian steroidogenesis and prostaglandin production in frog *Rana esculenta*, 123  
 Brechue, W.F. *see* Fukushima, S.  
 Brismar, H., Agrén, M. & Holtbäck, U.  $\beta$ -Adrenoceptor agonist sensitizes the dopamine-1 receptor in renal tubular cells, 333  
 Bruce, C.R. *see* Lee, J.S.  
 Byberg, L. *see* Hedman, A.  
 Cameron-Smith, D. *see* Lee, J.S.  
 Carré, F. *see* Lemoine, S.  
 Chennaoui, M., Gomez Merino, D., Lesage, J., Drogou, C. & Guezennec, C.Y. Effects of moderate and intensive training on the hypothalamo-pituitary-adrenal axis in rats, 113  
 Christensen, G. *see* Schiøtz Thorud, H.-M.  
 Christensen, G. *see* Woldbaek, P.R.  
 Christensen, S. *see* Jonassen, T.E.N.  
 Christova, T. *see* Kadinov, B.  
 Collins, D.E. *see* Yaspelkis III, B.B.  
 Crandall, C.G. *see* Kondo, N.  
 Delamarche, P. *see* Lemoine, S.  
 Diamant, M., Harms, M.P.M., Immink, R.V., Van Lieshout, J.J. & Van Montfrans, G.A. Twenty-four-hour non-invasive monitoring of systemic haemodynamics and cerebral blood flow velocity in healthy humans, 1  
 Drogou, C. *see* Chennaoui, M.  
 Durdanova, D. *see* Kadinov, B.  
 Eha, I. *see* Tähepöld, P.  
 Eintrei, C. *see* Björnström, H.C.  
 Ekblad, E. *see* Josefsson, M.  
 Elfström, P. *see* Tähepöld, P.  
 Ellingsen, O. *see* Schiøtz Thorud, H.-M.  
 Engberg, G. *see* Erhardt, S.  
 Erhardt, S. & Engberg, G. Increased phasic activity of dopaminergic neurones in the rat ventral tegmental area following pharmacologically elevated levels of endogenous kynurenic acid, 45  
 Fölmer, R. *see* Storr, M.  
 Fraga, S., Sampaio-Maia, B., Serrão, M.P. & Soares-da-Silva, P. Regulation of apical transporter of L-DOPA in human intestinal Caco-2 cells, 103  
 Fukushima, S., Abe, T., Shibayama, A. & Brechue, W.F. Comparison of viscoelastic characteristics in triceps surae between Black and White athletes, 183  
 Fukunaga, T. *see* Kubo, K.  
 Gagov, H. *see* Kadinov, B.  
 Gavin, T.P. & Wagner, P.D. Attenuation of the exercise-induced increase in skeletal muscle Flt-1 mRNA by nitric oxide synthase inhibition, 201  
 Gibson, K.J. *see* Westerlind, K.C.  
 Gobetti, A. *see* Bramucci, M.  
 Goloubeva, E.V. *see* Aleksandrova, N.P.  
 Gomes, P. & Soares-da-Silva, P. Dopamine-induced inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase activity requires integrity of actin cytoskeleton in opossum kidney cells, 93  
 Gomez Merino, D. *see* Chennaoui, M.  
 Granier, P. *see* Lemoine, S.  
 Grunditz, T. *see* Josefsson, M.  
 Guezennec, C.Y. *see* Chennaoui, M.  
 Harms, M.P.M. *see* Diamant, M.  
 Hassaf, D. *see* Arvola, L.  
 Hawley, J.A. *see* Lee, J.S.  
 Hedman, A., Byberg, L., Reneland, R. & Lithell, H.O. Muscle morphology, self-reported physical activity and insulin resistance syndrome, 325  
 Heide, A.M. *see* Jonassen, T.E.N.  
 Helgesen, K.G. *see* Arvola, L.  
 Hoen, I.B. *see* Woldbaek, P.R.  
 Holtbäck, U. *see* Brismar, H.  
 Horikawa, N. *see* Kondo, N.  
 Hügel, H. *see* Lee, J.S.  
 Ibarra, F.R., Jun Cheng, S.X., Agrén, M., Svensson, L.-B., Aizman, O. & Aperia, A. Intracellular sodium modulates the state of protein kinase C phosphorylation of rat proximal tubule Na<sup>+</sup>-K<sup>+</sup>-ATPase, 165  
 Illebekk, A. *see* Sjaastad, I.  
 Immink, R.V. *see* Diamant, M.  
 Inoue, Y. *see* Kondo, N.  
 Isaev, G.G. *see* Aleksandrova, N.P.  
 Itzev, D. *see* Kadinov, B.  
 Janjua, N.R. *see* Jonassen, T.E.N.  
 Jia, J. *see* Nyström, H.C.

- Johansson, M. *see* Nyström, H.C.
- Jonassen, T.E.N., Heide, A.M., Janjua, N.R. & Christensen, S. Collecting duct function in liver cirrhotic rats with early sodium retention, 237
- Josefsson, M., Grunditz, T., Ohlsson, T. & Ekblad, E. Sodium/iodide-symporter: distribution in different mammals and role in entero-thyroid circulation of iodide, 129
- Joumaa, W.H., Serrurier, B., Bigard, X. & Léoty, C. Nandrolone decanoate treatment induces changes in contractile responses of rat untrained fast-twitch skeletal muscle, 189
- Joyner, M.J. *see* Shastri, S.
- Jun Cheng, S.X. *see* Ibarra, F.R.
- Kadinov, B., Itzev, D., Gagov, H., Christova, Kolev, V., Duridanova, D. & Bolton, T.B. Induction of heme oxygenase in guinea-pig stomach: roles in contraction and in single muscle cell ionic currents, 297
- Kals, J. *see* Tähepõld, P.
- Kanchisa, H. *see* Kubo, K.
- Kiil, F. Mechanisms of intercellular hypertonicity and isotonic fluid absorption in proximal tubules of mammalian kidneys, 71
- Kiil, F. Mechanisms of transjunctional transport of NaCl and water in proximal tubules of mammalian kidneys, 55
- Kolev, V. *see* Kadinov, B.
- Kondo, N., Horikawa, N., Aoki, K., Shibasaki, M., Inoue, Y., Nishiyasu, T. & Crandall, C.G. Sweating responses to a sustained static exercise is dependent on thermal load in humans, 289
- Krisan, A.D. *see* Yaspelkis III, B.B.
- Kubo, K., Kanchisa, H. & Fukunaga, T. Effects of transient muscle contractions and stretching on the tendon structures *in vivo*, 157
- Kurjak, M. *see* Storr, M.
- Lambert, G. *see* Nyström, H.C.
- Lecoq, A.-M. *see* Nottin, S.
- Lee, J.S., Bruce, C.R., Tunstall, R.J., Cameron-Smith, D., Hügel, H. & Hawley, J.A. Interaction of exercise and diet on GLUT-4 protein and gene expression in Type I and Type II rat skeletal muscle, 37
- Lemoine, S., Granier, P., Tiffocche, C., Berthon, P.M., Thieulant, M.-L., Carré, F. & Delamarche, P. Effect of endurance training on oestrogen receptor alpha expression in different rat skeletal muscle type, 211
- Léoty, C. *see* Joumaa, W.H.
- Lesage, J. *see* Chennaoui, M.
- Lithell, H.O. *see* Hedman, A.
- Lunde, P.K. *see* Schiøtz Thorud, H.-M.
- Maccari, E. *see* Bramucci, M.
- McCarty, H.L. *see* Westerlind, K.C.
- Makhloufi, Y. *see* Mamchaoui, K.
- Mamchaoui, K., Makhloufi, Y. & Saumon, G. Glucose transporter gene expression in freshly isolated and cultured rat pneumocytes, 19
- Melnikov, A.L. *see* Arvola, L.
- Miano, A. *see* Bramucci, M.
- Murri, O. *see* Bramucci, M.
- N'Guyen, L.-D. *see* Nottin, S.
- Nishiyasu, T. *see* Kondo, N.
- Nottin, S., Vinet, A., Stecken, F., N'Guyen, L.-D., Ounissi, F., Lecoq, A.-M. & Obert, P. Central and peripheral cardiovascular adaptations to exercise in endurance-trained children, 85
- Nyström, H.C., Jia, J., Johansson, M., Lambert, G. & Bergström, G. Neurohormonal influences on maintenance and reversal of two-kidney one-clip renal hypertension, 245
- Obert, P. *see* Nottin, S.
- Ohlsson, T. *see* Josefsson, M.
- Ounissi, F. *see* Nottin, S.
- Persson, K. & Åhrén, B. The neuropeptide PACAP contributes to the glucagon response to insulin-induced hypoglycaemia in mice, 25
- Quassinti, L. *see* Bramucci, M.
- Reneland, R. *see* Hedman, A.
- Sampaio-Maia, B. *see* Fraga, S.
- Saumon, G. *see* Mamchaoui, K.
- Schiøtz Thorud, H.-M., Wisloff, U., Lunde, P.K., Christensen, G., Ellingsen, O. & Sejersted, O.M. Surgical manipulation, but not moderate exercise, is associated with increased cytokine mRNA expression in the rat soleus muscle, 219
- Schippert, A. *see* Björnström, K.
- Schusdziarra, V. *see* Storr, M.
- Sejersted, O.M. *see* Schiøtz Thorud, H.-M.
- Sejersted, O.M. *see* Sjaastad, I.
- Semb, S.O. *see* Sjaastad, I.
- Serrão, M.P. *see* Fraga, S.
- Serrurier, B. *see* Joumaa, W.H.
- Shastri, S., Toft, D.O. & Joyner, M.J. HSP70 and HSP90 expression in leucocytes after exercise in moderately trained humans, 139
- Shibasaki, M. *see* Kondo, N.
- Shibayama, A. *see* Fukushima, S.
- Singh, M.K. *see* Yaspelkis III, B.B.
- Sjaastad, I., Bentzen, J.G., Semb, S.O., Ilebakk, A. & Sejersted, O.M. Reduced calcium tolerance in rat cardiomyocytes after myocardial infarction, 261
- Sjölinder, A. *see* Björnström, K.
- Sjöquist, P.-O. *see* Wang, Q.-D.
- Soares-da-Silva, P. *see* Fraga, S.
- Soares-da-Silva, P. *see* Gomes, P.
- Starkopf, J. *see* Tähepõld, P.
- Stecken, F. *see* Nottin, S.
- Storr, M., Fölmer, R., Kurjak, M., Schusdziarra, V. & Allescher, H.D. Okadaic acid inhibits relaxant neural transmission in rat gastric fundus *in vitro*, 29
- Strange, R. *see* Westerlind, K.C.
- Svensson, L.-B. *see* Ibarra, F.R.
- Taal, G. *see* Tähepõld, P.
- Tähepõld, P., Elfström, P., Eha, I., Kals, J., Taal, G., Talonpoika, A., Valen, G., Vaage, J. & Starkopf, J. Exposure of rats to hyperoxia enhances relaxation of isolated aortic rings and reduces infarct size of isolated hearts, 271
- Talonpoika, A. *see* Tähepõld, P.

- Thiculant, M.-L. *see* Lemoine, S.  
 Thorén, P. *see* Wang, Q.-D.  
 Tiffiche, C. *see* Lemoine, S.  
 Toft, D.O. *see* Shastri, S.  
 Tokuno, S. *see* Wang, Q.-D.  
 Tønnessen, T. *see* Woldback, P.R.  
 Trevino, B. *see* Yaspelkis III, B.B.  
 Tunstall, R.J. *see* Lee, J.S.
- Vaage, J. *see* Tähepõld, P.  
 Valen, G. *see* Tähepõld, P.  
 Valen, G. *see* Wang, Q.-D.  
 Van Lieshout, J.J. *see* Diamant, M.  
 Van Montfrans, G.A. *see* Diamant, M.  
 Vinet, A. *see* Nottin, S.
- Wagner, P.D. *see* Gavin, T.P.  
 Wang, Q.-D., Tokuno, S., Valen, G., Sjöquist, P.-O. & Thorén, P. Cyclic fluctuations in the cardiac performance of the isolated Langendorff-perfused mouse heart: pyruvate abolishes the fluctuations and has an anti-ischaemic effect, 279
- Westerlind, K.C., McCarty, H.L., Gibson, K.J. & Strange, R. Effect of exercise on the rat mammary gland: implications for carcinogenesis, 147  
 Wisloff, U. *see* Schiøtz Thorud, H.-M.  
 Woldback, P.R., Hoen, I.B., Christensen, G. & Tønnessen, T. Gene expression of colony-stimulating factors and stem cell factor after myocardial infarction in the mouse, 173
- Yaspelkis III, B.B., Singh, M.K., Trevino, B., Krisan, A.D. & Collins, D.E. Resistance training increases glucose uptake and transport in rat skeletal muscle, 315  
 Ytrehus, K. *see* Arvola, L.
- Zerani, M. *see* Bramucci, M.



## Subject index

- A10, 45  
Achizophrenia, 45  
action potential duration, 11  
adenosine, 11  
 $\alpha$ -adrenergic agonists, 165  
adrenocorticotropin hormone (ACTH), 113  
aerobic exercise training, 315  
aerobic training, 85  
airway receptors, 253  
alveolar epithelium, 19  
alveolar liquid clearance, 19  
apoptosis, 147  
amiloride, 237  
anaesthetic mechanisms, 227  
aquaporins, 71
- blood lipids, 325  
blood pressure, 1, 245  
breast cancer, 147  
burst activity, 45
- Caco-2 cells, 103  
capillary supply, 325  
cardiac output, 1, 85  
cerebral blood flow velocity, 1  
children, 85  
cirrhosis, 237  
citrate synthase, 315  
collecting duct, 237  
contractile force, 11  
contraction, 297  
corticosterone, 113  
corticotropin-releasing factor (CRF), 113  
cutaneous vascular response, 289  
cytokine, 219  
cytokines, 173  
cytoskeleton, 93
- 2,6-di-isopropylphenol, 227  
D-cycloserine, 45  
differentiation, 19  
diffusion coefficient, 55  
D-NAME, 201  
Donnan equilibrium, 55  
dopamine, 45, 93, 165  
Doppler, 1  
down-regulation, 139
- 17 $\beta$ -estradiol, 123  
effective osmolality, 55  
entro-thyroid circulation, 129  
enzyme activity, 297  
estradiol, 147  
excitation-contraction coupling, 189  
exercise, 85, 219  
exercise training, 37, 147
- fetal liver kinase-1 receptor, 201  
fluctuation, 279  
fms-like tyrosine kinase-1 receptor, 201  
free vibration technique, 183
- gastric mucosa, 129  
gene expression, 173  
glucagon secretion, 25  
GLUT4, 315  
glycogen, 315  
G-protein-coupled receptors, 333  
guinea-pig, 11, 297
- haemin, 297  
heart, 279  
heart failure, 173, 261  
heme oxygenase, 297  
heterologous sensitization, 333  
hexokinase, 315  
high-fat diet, 37  
histochemistry, 297  
human, 157  
hyperoxia, 271  
hypertension, 245  
hypoglycaemia, 25  
hysteresis, 157
- insulin resistance syndrome, 325  
insulin-resistance, 37  
intact and skinned fibres, 189  
intra/extracellular ions, 261  
intracellular sodium, 165  
intravenous, 227  
iodide transport, 129  
ion exchangers, 261  
ionic currents, 297  
ischaemia-reperfusion of the heart, 271  
isolated aortic rings, 271  
isometric handgrip exercise, 289  
isoproterenol, 11, 333
- K<sup>+</sup>-ATPase, 165  
K<sup>+</sup> contracture, 189  
kidney, 245  
kidney medulla, 245  
kynurenic acid, 45
- lateral intercellular space, 71  
L-DOPA, 103  
left ventricular dimensions, 85  
L-NAME, 201  
load compensation, 253  
L-type amino acid transporter, 103
- mannitol, 55  
maximal oxygen consumption, 139

- mice, 25  
middle cerebral artery, 1  
MK 801, 45  
molecular chaperones, 139  
multiprobe RNase protection assay, 219  
murine, 173  
muscle fibre composition, 325  
muscle triacylglycerol, 37  
muscle type, 211  
muscle-tendon complex, 183  
myocardial infarction, 173, 261, 271
- $\text{Na}^+$ , 165  
Na/Ca-exchanger, 261  
 $\text{Na}^+/\text{K}^+$ -ATPase, 93, 333  
Na-K-ATPase, 237  
NANC relaxation, 29  
neurone, 227  
neutral endopeptidase 2411, 123  
NIS, 129  
NO, 29  
non-thermal factors, 289
- oestrogen receptor  $\alpha$ , 211  
OK cells, 93  
okadaic acid, 29
- PACAP, 25  
papillary muscle, 11  
perfusion, 279  
physical activity, 325  
physical training adaptation, 113  
PNU 156561A, 45  
progesterone, 147  
proliferation, 147  
propofol, 227  
prostaglandin  $\text{E}_2$ , 123  
protein kinase C, 165  
protein kinases, 11  
protein phosphatase 1 and 2A, 29  
proximal tubular NaCl transport, 71  
proximal tubular  $\text{S}_1$  and  $\text{S}_2$  segments, 55  
pyruvate mouse, 279
- race difference, 183  
rat, 219, 227  
rat model, 147  
rats, 245  
receptor recruitment, 333  
reflection coefficient, 55, 71  
renin-angiotensin system, 245  
resistive load, 253  
restraint stress, 113  
RT-PCR, 211
- salivary glands, 129  
signal transduction, 11  
skeletal muscle, 211, 219  
skeletal muscles, 189  
smooth muscle, 29, 297  
sodium retention, 237  
sodium transport, 19  
stiffness, 157  
stress, 139  
sympathetic nervous system, 245
- tendon, 157  
terminal end bud, 147  
thermal factors, 289  
thyroid gland, 129  
tight junction, 55, 71  
training, 85, 211  
twitch, 189  
tyrosine kinase, 227
- ultrasonography, 157  
unclipping, 245  
upper airway muscles, 253  
uptake, 103
- vascular endothelial growth factor (VEGF), 201  
vascular function, 271  
vessel reactivity, 271  
visco-elasticity, 183

